Brain Research, 446 (1988) 407–409
Elsevier

Author Index

Agid, Y., see Maloteaux, J.-M., 321
Akeyson, E.W., see Knuepfer, M.M., 17
Amiri, I., see Ungerer, A., 205
Anton, F., see Szolcsanyi, J., 262
Aou, S., see Koyama, Y., 199
Araki, H., see Ishihara, A., 195
Azizi, S.A., see Cheng, J.-T., 90

Baba, S., see Sakaue, M., 343
Barnhill, J.G., see Miller, L.G., 314
Besson, J.M., see Rivot, J.P., 333
Bidard, J.-N., see Mourre, C., 106
Bol, J.G.J.M., see De Vente,, J., 387
Boulanger, Y., see Ungerer, A., 205

Caldwell, J.D., Jirikowski, G.F., Greer, E.R., Stumpf, W.E. and Pedersen, C.A., Ovarian steroids and sexual interaction alter oxytocinergic content and distribution in the basal forebrain, 236
Campochiaro, P.A., see Friedman, Z., 11
Cannon-Spoor, H.E., see Freed, W.J., 133
Caprio, J., see Marui, T., 178
Casanova, C., see Molotchnikoff, S., 67
Cérat, A., see Molotchnikoff, S., 67
Chapin, J.K., see Cheng, J.-T., 90
Cheng, J.-T., see Sessler, F.M., 77
Cheng, J.-T., Sessler, F.M., Azizi, S.A., Chapin, J.K. and Waterhouse, B.D., Electrophysiological actions of norepinephrine in rat lateral hypothalamus. II. An in vitro study of the effects of iontophoretically applied norepinephrine on LH neuronal responses to γ-aminobutyric acid (GABA), 90
Choi, D.W., see Koh, J.-Y., 374
Choi, D.W., see Monyer, H., 144
Clough, R.W., Hoffman, G.E. and Sladek, C.D., Peripubertal development of noradrenergic stimulation of luteinizing hormone-releasing hormone neurosecretion in vitro, 121
Coceani, F., Lees, J. and Dinarello, C.A., Occurrence of interleukin-1 in cerebrospinal fluid of the conscious rat, 245
Coupet, J., see Hinson, J., 379
Czéh, G., Obih, J.-C.A. and Somjen, G.G., The effect of changing extracellular potassium concentration on synaptic transmission in isolated spinal cords, 50

Davis, K.D. and Dostrovsky, J.O., Cerebrovascular application of bradykinin excites central sensory neurons, 401
De Barry, J., see Ungerer, A., 205
Deckert, J. and Jorgensen, M.B., Evidence for pre- and postsynaptic localization of adenosine A1 receptors in the CA1 region of rat hippocampus: a quantitative autoradiographic study, 161
DeLanney, L.E., see Ricaurte, G.A., 165
Del Fiacco, M., Perra, M.T., Quartu, M., Rosa, M.D., Zucca, G. and Levanti, M.C., Evidence for the presence of substance P-like immunoreactivity in the human cerebellum, 173
De Vente, J., Bol, J.G.J.M., Hudson, L., Schipper, J. and Steinbusch, H.W.M., Atrial natriuretic factor-responding and cyclic guanosine monophosphate (cGMP)-producing cells in the rat hippocampus: a combined micropharmacological and immunocytochemical approach, 387
De Volder, A., see Wanet-Defalque, M.-C., 369
Devor, M. and Dubner, R., Centrifugal activity in afferent C-fibers influences the spontaneous afferent barrage generated in nerve-end neuromas, 396
Dinarello, C.A., see Coceani, F., 245
Dooms, G., see Wanet-Defalque, M.-C., 369
Dostrovsky, J.O., see Davis, K.D., 401
Dubner, R., see Devor, M., 396

Eckenstein, F., Transient expression of NGF-receptor-like immunoreactivity in postnatal rat brain and spinal cord, 149
Emson, P.C., see Kuwayama, Y., 219

Feer, H., see Rudolph, K., 159
Forman, D.S., see Smith, R.S., 26
Frederickson, C.J., Hernandez, M.D., Goik, S.A., Morton, J.D. and McGinty, J., Loss of zinc staining from hippocampal mossy fibers during kainic acid induced seizures: a histofluorescent study, 383
Freed, W.J. and Cannon-Spoor, H.E., Cortical lesions increase reinnervation of the dorsal striatum by substantia nigra grafts, 133
Friedl, W., see Hebebrand, J., 251
Friedman, Z., Hackett, S.F. and Campochiaro, P.A., Human retinal pigment epithelial cells possess muscarinic receptors coupled to calcium mobilization, 11
Fujita, I., see Koyama, Y., 199
Fukuda, J., see Hirono, T., 189
Fukuda, Y., see Sumitomo, I., 354

Gebhart, G.F., see Ren, K., 285
Giordano, M., Hagenmeyer-Houser, S.H. and Sanberg, P.R., Intraparenchymal fetal striatal transplants and recovery in kainic acid lesioned rats, 183
Goffinet, A., see Wanet-Defalque, M.-C., 369
Goik, S.A., see Frederickson, C.J., 383
Gonzalez, H.A., see Wang, P.S., 363
Greenblatt, D.J., see Miller, L.G., 314
Greer, E.R., see Caldwell, J.D., 236

Hackett, S.F., see Friedman, Z., 11
Hagenmeyer-Houser, S.H., see Giordano, M., 183
Handwerker, H.O., see Szolcsanyi, J., 262
Hayaishi, O., see Suzuki, S., 1
Hebebrand, J., Friedl, W., Reichelt, R., Schmitz, E., Möller, P. and Propping, P., The shark GABA–benzodiazepine receptor: further evidence for a not so late phylogenetic appearance of the benzodiazepine receptor, 251
Hernandez, M.D., see Frederickson, C.J., 383
Hinson, J., Rauh, C. and Coupet, J., Neuropeptide Y stimulates inositol phospholipid hydrolysis in rat brain miniprisms, 379
Hirono, T., Torimitsu, K., Kawana, A. and Fukuda, J., Recognition of artificial microstructures by sensory nerve fibers

in culture, 189
Hirose, Y., see Kurasawa, I., 113
Hoffman, G.E., see Clough, R.W., 121
Hsiao, C.-F., see Sumitomo, I., 354
Hudson, L., see De Vente, J., 387

Introini-Collison,, I.B., see McGaugh, J.L., 37
Irwin, I., see Ricaurte, G.A., 165
Ishihara, A., Naitoh, H., Araki, H. and Nishihira, Y., Soma size and oxidative enzyme activity of motoneurones supplying the fast twitch and slow twitch muscles in the rat, 195

Javoy-Agid, F., see Maloteaux, J.-M., 321
Jirikowski, G.F., see Caldwell, J.D., 236
Jorgensen, M.B., see Deckert, J., 161

Kagamiyama, H., see Suzuki, S., 1
Kanda, K., see Sato, Y., 225
Kasahara, Y., see Marui, T., 178
Kawana, A., see Hirono, T., 189
Kawasaki, T., see Sato, Y., 225
Kiyohara, S., see Marui,T., 178
Knuepfer, M.M., Akeyson, E.W. and Schramm, L.P., Spinal projections of renal afferent nerves in the rat, 17
Koh, J.-Y. and Choi, D.W., Cultured striatal neurons containing NADPH-diaphorase or acetylcholinesterase are selectively resistant to injury by NMDA receptor agonists, 374
Koyama, Y., Fujita, I., Aou, S. and Oomura, Y., Proceptive presenting elicited by electrical stimulation of the female monkey hypothalamus, 199
Krăuchi, K., see Rudolph, K., 159
Krieger, E.M., see Menani, J.V., 295
Kurasawa, I., Hirose, Y., Sunada, T. and Nakamura, Y., Phase-linked modulation of excitability of presynaptic terminals of low-threshold afferent fibers in the inferior alveolar nerve during cortically induced fictive mastication in the guinea pig, 113
Kuwayama, Y., Emson, P.C. and Stone, R.A., Pterygopalatine ganglion cells contain neuropeptide Y, 219

Laduron, P.-M., see Maloteaux, J.-M., 321
Lamour, Y., see Rascol, O., 303
Langston, J.W., see Ricaurte, G.A., 165
Laterre, E.C., see Maloteaux, J.-M., 321
Lawrence, D., see Lawrence, J.A.M., 212
Lawrence, J.A.M., Poulin, P., Lawrence, D. and Lederis, K., [^{3}H]Arginine vasopressin binding to rat brain: a homogenate and autoradiographic study, 212
Lazdunski, M., see Mourre, C., 106
Lederis, K., see Lawrence, J.A.M., 212
Lees, J., see Coceani, F., 245
Levanti, M.C., see Del Fiacco, M., 173
Luabeya, M.A.K., see Maloteaux, J.-M., 321

Machado, B.H., see Menani, J.V., 295
Maloteaux, J.-M., Luabeya, M.A.K., Vanisberg, M.-A., Laterre, E.C., Laduron, P.-M., Javoy-Agid, F. and Agid, Y., Benzodiazepine receptors in normal human brain, in Parkinson's disease and in progressive supranuclear palsy, 321
Marui, T., Caprio, J., Kiyohara, S. and Kasahara, Y., Topographical organization of taste and tactile neurons in the facial lobe of the sea catfish, *Plotosus lineatus*, 178
McGaugh, J.L., Introini-Collison, I.B. and Nagahara, A.H., Memory-enhancing effects of posttraining naloxone: involvement of β-noradrenergic influences in the amygdaloid complex, 37
McGinty, J., see Frederickson, C.J., 383
Mélan, C., see Ungerer, A., 205
Menani, J.V., Machado, B.H., Krieger, E.M. and Salgado, H.C., Tachycardia during the onset of one-kidney, one-clip renal hypertension: role of the renin–angiotensin system and AV3V tissue, 295
Metz, R., see Wanet-Defalque, M.-C., 369
Michel, C., see Wanet-Defalque, M.-C., 369
Miletic, V. and Tan, H., Iontophoretic application of calcitonin gene-related peptide produces a slow and prolonged excitation of neurons in the cat lumbar dorsal horn, 169
Miller, L.G., Greenblatt, D.J., Barnhill, J.G., Thompson, M.L. and Shader, R.I., Modulation of benzodiazepine receptor binding in mouse brain by adrenalectomy and steroid replacement, 314
Möller, P., see Hebebrand, J., 251
Molotchnikoff, S., Cérat, A. and Casanova, C., Influences of cortico-pretectal fibers on responses of rat pretectal neurons, 67
Monyer, H. and Choi, D.W., Morphinans attenuate cortical neuronal injury induced by glucose deprivation in vitro, 144
Morton, J.D., see Frederickson, C.J., 383
Mourre, C., Bidard, J.-N. and Lazdunski, M., High affinity receptors for the bee venom MCD peptide. Quantitative autoradiographic localization at different stages of brain development and relationship with MCD neurotoxicity, 106

Nagahara, A.H., see McGaugh, J.L., 37
Naitoh, H.,see Ishihara, A., 195
Nakamura, Y., see Kurasawa, I., 113
Nishihira, Y., see Ishihara, A., 195

Oberg, P.A., see Rechthand, E., 61
Obih, J.-C.A., see Czéh, G., 50
Oomura, Y., see Koyama, Y., 199

Pedersen, C.A., see Caldwell, J.D., 236
Perra, M.T., see Del Fiacco, M., 173
Pointis, D., see Rivot, J.P., 333
Porter, J.C., see Wang, P.S., 363
Poulin, P., see Lawrence, J.A.M., 212
Propping, P., see Hebebrand, J., 251

Quartu, M., see Del Fiacco, M., 173

Randich, A., see Ren, K., 285
Rapoport, S.I., see Rechthand, E., 61
Rascol, O. and Lamour, Y., Differential effects of M_1 and M_2 muscarinic drugs on septohippocampal, hippocampal and cortical neurons in the rat, 303
Rauh, C., see Hinson, J., 379
Rechthand, E., Sato, S., Oberg, P.A. and Rapoport, S.I., Sciatic nerve blood flow response to carbon dioxide, 61
Reeh, P.W., see Szolcsanyi, J., 262
Reichelt, R., see Hebebrand, J., 251
Reinbolt, J., see Ungerer, A., 205
Ren, K., Randich, A. and Gebhart, G.F., Vagal afferent modulation of a nociceptive reflex in rats: involvement of spinal opioid and monoamine receptors, 285
Ricaurte, G.A., DeLanney, L.E., Irwin, I. and Langston, J.W., Toxic effects of MDMA on central serotonergic neu-

rons in the primate: importance of route and frequency of drug administration, 165
Rivot, J.P., Pointis, D. and Besson, J.M., Morphine increases 5-HT metabolism in the nucleus raphe magnus: an in vivo study in freely moving rats using 5-hydroxyindole electrochemical detection, 333
Roeske, W.R., see Wang, J.-x., 155
Rosa, M.D., see Del Fiacco, M., 173
Rudolph, K., Wirz-Justice, A., Kräuchi, K. and Feer, H., Static magnetic fields decrease nocturnal spinal cAMP in the rat, 159

Saito, N., see Sakaue, M., 343
Sakaue, M., Saito, N., Taniguchi, H., Baba, S. and Tanaka, C., Immunohistochemical localization of γ-aminobutyric acid in the rat pituitary gland and related hypothalamic regions, 343
Salgado, H.C., see Menani, J.V., 295
Sanberg, P.R., see Giordano, M., 183
Sato, S., see Rechthand, E., 61
Sato, Y., Kanda, K. and Kawasaki, T., Target neurons of flocular middle zone inhibition in medial vestibular nucleus, 225
Schipper, J., see De Vente, J., 387
Schmidt, J.T. and Shashoua, V.E., Antibodies to ependymin block the sharpening of the regenerating retinotectal projection in goldfish, 269
Schmitz, E., see Hebebrand, J., 251
Schmitz-Bourgeois, M., see Ungerer, A., 205
Schramm, L.P., see Knuepfer, M.M., 17
Sessler, F.M., Cheng, J.-T. and Waterhouse, B.D, Electrophysiological actions of norepinephrine in rat lateral hypothalamus.I. Norepinephrine-induced modulation of LH neuronal responsiveness to afferent synaptic inputs and putative neurotransmitters, 77
Sessler, F.M., see Cheng, J.-T., 90
Shader, R.I., see Miller, L.G., 314
Shashoua, V.E., see Schmidt, J.T., 269
Sladek, C.D., see Clough, R.W., 121
Smith, R.S. and Forman, D.S., Organelle dynamics in lobster axons: anterograde and retrograde particulate organelles, 26
Somjen, G.G., see Czéh, G., 50
Steinbusch, H.W.M., see De Vente, J., 387
Stone, R.A., see Kuwayama, Y., 219
Stumpf, W.E., see Caldwell, J.D., 236
Sumitomo, I., Hsiao, C.-F. and Fukuda, Y., Two types of thalamic reticular cells in relation to the two visual thalamocortical systems in the rat, 354
Sunada, T., see Kurasawa, I., 113
Suzuki, S., Watanabe, Y., Tsubokura, S., Kagamiyama, H. and Hayaishi, O., Decrease in tetrahydrobiopterin content and neurotransmitter amine biosynthesis in rat brain by an inhibitor of guanosine triphosphate cyclohydrolase, 1
Szolcsanyi, J., Anton, F., Reeh, P.W. and Handwerker, H.O., Selective excitation by capsaicin of mechano-heat sensitive nociceptors in rat skin, 262

Tan, H., see Miletic, V., 169
Tanaka, C., see Sakaue, M., 343
Taniguchi, H., see Sakaue, M., 343
Thompson, M.L., see Miller, L.G., 314
Torimitsu, K., see Hirono, T., 189
Tsubokura, S., see Suzuki, S., 1

Ungerer, A., Schmitz-Bourgeois, M., Mélan, C., Boulanger, Y., Reinbolt, J., Amiri, I. and De Barry, J., γ-L-Glutamyl-L-aspartate induces specific deficits in long-term memory and inhibits [^{3}H]glutamate binding on hippocampal membranes, 205

Vanisberg, M.-A., see Maloteaux, J.-M., 321
Veraart, C., see Wanet-Defalque, M.-C., 369

Wanet-Defalque, M.-C., Veraart, C., De Volder, A., Metz, R., Michel, C., Dooms, G. and Goffinet, A., High metabolic activity in the visual cortex of early blind human subjects, 369
Wang, J.-x., Roeske, W.R., Wang, W. and Yamamura, H.I., Himbacine recognizes a high affinity subtype of M_2 muscarinic cholinergic receptors in the rat cerebral cortex, 155
Wang, P.S., Gonzalez, H.A. and Porter, J.C., Saline ingestion stimulates the in situ molar activity of tyrosine hydroxylase in the median eminence and superior cervical ganglion, 363
Wang, W., see Wang, J.-x., 155
Watanabe, Y., see Suzuki, S., 1
Waterhouse, B.D., see Cheng, J.-T., 90
Waterhouse, B.D., see Sessler, F.M., 77
Wirz-Justice, A., see Rudolph, K., 159

Yamamura, H.I., see Wang, J.-x., 155

Zucca, G., see Del Fiacco, M., 173